

Appl. No. 10/061,727
Amdt. dated December 17, 2004
Resp. to Advisory Action dated Sept. 9, 2004

CLAIMS

This listing of claims will replace all prior versions and listings of claims in the application:

1. (Original) An isolated polynucleotide comprising SEQ ID NO:1 wherein the nucleic acid at 1792 is A or C.
2. (Previously presented) An isolated polynucleotide comprising a nucleic acid that encodes a polypeptide comprising SEQ ID NO:2, wherein the amino acid 598 is Thr or Pro.
3. (Canceled)
4. (Canceled)
5. (Currently amended) An isolated polynucleotide comprising a molecule selected from the group consisting of:
 - a) A polynucleotide that encodes a polypeptide comprising amino acid residues 384-687 of SEQ ID NO:2, wherein the amino acid at 598 is Thr or Pro;
 - b) A polynucleotide that encodes a polypeptide comprising amino acid residues 379-687 of SEQ ID NO:2, wherein the amino acid at 598 is Thr or Pro;
 - ~~c) A polynucleotide that encodes a polypeptide comprising amino acid residues 389-685 of SEQ ID NO:4;~~
 - ~~d) A polynucleotide that encodes a polypeptide comprising amino acid residues 379-685 of SEQ ID NO:4;~~
 - e)c) A polynucleotide that encodes a polypeptide comprising amino acid residues 449-687 of SEQ ID NO:2, wherein the amino acid at 598 is Pro or Thr;
 - ~~f) A polynucleotide that encodes a polypeptide comprising amino acid residues 449-685 of SEQ ID NO:4;~~
 - g)d) A polynucleotide that encodes a fragment of a polypeptide described in (a-[[f]]c), wherein the fragment interacts with an IL-1R signal transduction factor;
 - h)e) An isolated nucleic acid molecule that hybridizes to either strand of a denatured, double-stranded DNA comprising the polynucleotide of any one of [[e-g]]c under conditions of moderate stringency in 50% formamide and 6XSSC, at 42°C with washing conditions of 60°C, 0.5XSSC, 0.1% SDS; wherein the isolated nucleic acid encodes a polypeptide that interacts with an IL-1R signal transduction factor;

Appl. No. 10/061,727
Amdt. dated December 17, 2004
Resp. to Advisory Action dated Sept. 9, 2004

- i)f) An isolated nucleic acid molecule that encodes a polypeptide that is at least 85% identical to the polypeptides described in ~~a)-g)~~a), b), c), wherein the polypeptide interacts with an IL-1R signal transduction factor;
- j)g) A polynucleotide that is degenerate to any of the polynucleotides of ~~a)-i)~~a), b), c), e), f).
6. (Original) An expression vector comprising a polynucleotide of claim 5.
7. (Original) An expression vector comprising a polynucleotide that encodes a polypeptide comprising SEQ ID NO:2, wherein the amino acid residue at 598 is Pro or Thr.
8. (Canceled)
9. (Previously presented) A host cell comprising the vector of claim 6.
10. (Original) A process of preparing a polypeptide, the process comprising culturing a host cell of claim 9 under conditions promoting expression of the polypeptide.
11. (Original) A process of preparing a polypeptide, the process comprising culturing a host cell transformed with a vector of claim 7 under conditions promoting expression of the polypeptide.
12. (Canceled)
13. (Canceled)
14. (Canceled)